

2.5%BROILER PREMIX

ANALYSIS	(0-18 Days)	(18 days to end)
Moisture max %	9,00	9,00
Crude protein %	22,00	20,20
Crude fat %	1,24	1,20
Crude ash %	45,00	43,00
Crude fiber %	1,50	1,80
Calcium %	19,00	19,00
Total Phosphorus min %	5,70	5,00
Av. Phosphorus %	9,60	8,80
Sodium %	0,04	0,06
Chlorine%	0,58	0,45
Total Lysine min%	3,20	2,50
Total Methionine %	8,10	6,68
Total Met + Cys.min %	8,65	6,80
Total Threonine min %	1,20	1,10
Total Tryptophan min %	0,20	0,23
Total Arginine min %	0,76	0,75
Total Valine min %	0,40	0,42
ME Energy Kcal	1200	1170

VITAMINS

Vitamin A IU	440 000	400 000
Vitamin D3 IU	96 000	80 000
Vitamin E (mg)	1000	700
Vitamin K3 (mg)	80	80
Vitamin B1 (mg)	45	40
Vitamin B2 (mg)	80	80
Vitamin B3 (mg)	400	400
Vitamin B6 (mg)	40	60
Vitamin B12 (mg)	0,8	0,8
Folic Acid (mg)	40	40
Nicotinic Acid (Niacin) (mg)	600	600
Choline Chloride (mg)	12 000	14 000
Biotin (mg)	4	4

SIMYA Feed Additives Ltd

Adres: Aziziye Mahallesi Kulođlu Sokak 15/4 ÇANKAYA ANKARA/TURKEY
 Tel: +90 312 440 81 51 Fax: +90 312 440 81 75

ANALYSIS	(0-18 Days)	(18 days to end)
Iron	900	900
Copper	300	300
Manganese	1.200	1.200
Zinc	1.400	1.400
Iodium	40	40
Selenium	8	8
OTHER ADDITIVES		
Antioxydant Blend (mg)	2.000	2.000
Phytase FTU/kg	20.000	20.000
E. 1,4 β Xylanase (TXT/kg)	22.400	22.400
E. 1,4 β Glucanase (TXT/kg)	10.000	10.000
Ethoxyquin (mg/kg)	100	100
Mannan (mg/kg)	21.600	21.600
Glucan (mg/kg)	8.800	8.800
Organic Acid (mg/kg)	20.000	20.000

***SIMYA BROILER 2,5% PREMIX** is prepared by using satisfied formulation and mixing technology according to the regional nutritional requirement, seasonal conditions, target feed specifications and performances. SIMYA 2,5% BROILER PREMIX is added 25 kg into the each metric ton feed.